

Angola

Total number of states \ provinces: 18

Immunization

Immunization coverage (2009 nat. WHO-UNICEF estimates) DTP3 73% Pol3 73%

Non Polio AFP Profile

<i>Percent 0-dose (national)</i> 12.7%	<i>Percent states with < 10% 0-dose NPAFP children</i> Non-adjusted 66.7% (2 of 3) % within acceptable range 100% (3 of 3)
<i>Percent 4+ doses (national)</i> 28.7%	<i>Percent states with > 80% 4+dose NPAFP children</i> Non-adjusted 0% (0 of 3) % within acceptable range 0% (0 of 3)

Independent Monitoring (IM data were not collected, or were not available in detail for all rounds shown)

<i>Campaign type</i> (mixed rounds abbreviated as: t, b, 1, 3)	<i>Start date</i>	<i>National indicator</i>		<i>Sub national indicator</i>	
		Percent missed children (finger marking)		Percent of states with < 10% missed children	
		House to House (%)	Out of house (%)	House to House	Out of house
SNID-mOPV1	04-Jun-10	9.6	9.5	50% (1 of 2)	50% (1 of 2)
NID-tOPV, 1	11-Jun-10	12.4	14.9	42% (5 of 12)	67% (8 of 12)
SNID-mOPV1	18-Jun-10	6.5	6.8	100% (1 of 1)	100% (1 of 1)
NID-bOPV	06-Aug-10	12.5	13.9	54% (7 of 13)	50% (6 of 12)
NID-bOPV	13-Aug-10	16.9	17.4	50% (1 of 2)	50% (1 of 2)
NID-bOPV	01-Oct-10	7.6	9.7	88% (15 of 17)	82% (14 of 17)
NID-mOPV1	29-Oct-10	9	9.9	72% (13 of 18)	67% (12 of 18)
SNID-OPVm1	10-Dec-10				
SNID-OPVm1	26-Dec-10				
SNID-mOPV1	25-Feb-11	6.7	7.5	100% (6 of 6)	83% (5 of 6)
NID-bOPV, 1	25-Mar-11	7.3	8.4	72% (13 of 18)	67% (12 of 18)
NID-bOPV, 1	28-Apr-11	6.1	8	100% (15 of 15)	87% (13 of 15)
NID-OPVb	27-May-11				

Surveillance

Polio Cases (W1, W3, and VDPVs)

<i>Serotype</i>	<i>Count</i>				<i>Date of onset for last case</i>
	2010 (Jan-Dec)		2011 (Jan-present)		
	cases	districts	cases	districts	
W1	33	18	4	1	27-Mar-11

Non Polio AFP data

<i>NPAFP rate (national)</i> 2.8	<i>Percent states with NPAFP rate >= 2</i> Non-adjusted 94.4% (17 of 18) % within acceptable range 100% (18 of 18)
<i>% adequate stools (nat.)</i> 84.3%	<i>Percent states with adequate stool proportion >= 80%</i> Non-adjusted 83.3% (15 of 18) % within acceptable range 94.4% (17 of 18)
<i>Percent of population <15 years of age living in states / prov. where both surveillance indicators are met</i> 65.8%	

Poliovirus History

The WPV1 viruses in Angola during the second half of 2010 and 2011 represented a continuing outbreak resulting from importation from India in 2007. Only one genetic cluster has been observed. In 2010, cases were distributed more widely including the northeastern part of the country and included moderate regional transmission. The cases in 2011 represented continued circulation from 2010 and not a new importation and marked the fourth consecutive year of transmission of this virus. Based upon multiple isolates from 2010 with less genetic linkage than expected, there is indication that surveillance may have gaps at the sub-national level.

Angola

Total number of states \ provinces: 18

Data Completeness

Percent of NPAFP cases with unknown age

0%

Percent of 6-35 month old NPAFP cases with unknown dose history

10.7%