

Angola

Total number of states \ provinces: 18

Immunization

Immunization coverage (2009 nat. WHO-UNICEF estimates) DTP3 91% Pol3 92%

Non Polio AFP Profile

<i>Percent 0-dose (national)</i> 13.7%	<i>Percent states with < 10% 0-dose NPAFP children</i> Non-adjusted 25% (1 of 4) % within acceptable range 75% (3 of 4)
<i>Percent 4+ doses (national)</i> 30.1%	<i>Percent states with > 80% 4+dose NPAFP children</i> Non-adjusted 0% (0 of 4) % within acceptable range 0% (0 of 4)

Independent Monitoring (IM data were not collected, or were not available in detail for all rounds shown)

<i>Campaign type</i> (mixed rounds abbreviated as: t, b, 1, 3)	<i>Start date</i>	<i>National indicator</i>		<i>Sub national indicator</i>	
		Percent missed children (finger marking)		Percent of states with < 10% missed children	
		House to House (%)	Out of house (%)	House to House	Out of house
NID-bOPV	01-Oct-10	7.6	9.7	88% (15 of 17)	82% (14 of 17)
NID-mOPV1	29-Oct-10	9	9.9	72% (13 of 18)	67% (12 of 18)
SNID-OPVm1	03-Dec-10				
SNID-OPVm1	26-Dec-10				
SNID-mOPV1	25-Feb-11	6.7	7.5	100% (6 of 6)	83% (5 of 6)
NID-bOPV+m1	25-Mar-11	7.3	8.4	72% (13 of 18)	67% (12 of 18)
NID-bOPV+m1	28-Apr-11	6.1	8	100% (15 of 15)	87% (13 of 15)
NID-bOPV+m1	27-May-11	6.7	7.7	93% (14 of 15)	87% (13 of 15)
SNID-bOPV	08-Jul-11	8.6	8	75% (3 of 4)	75% (3 of 4)

Surveillance

Polio Cases (W1, W3, and VDPVs)

<i>Serotype</i>	Count				Date of onset for last case
	2010 (Jan-Dec)		2011 (Jan-present)		
	cases	districts	cases	districts	
W1	33	18	4	1	27-Mar-11

Non Polio AFP data

<i>NPAFP rate (national)</i> 2.7	<i>Percent states with NPAFP rate >= 2</i> Non-adjusted 83.3% (15 of 18) % within acceptable range 100% (18 of 18)
<i>% adequate stools (nat.)</i> 85.9%	<i>Percent states with adequate stool proportion >= 80%</i> Non-adjusted 77.8% (14 of 18) % within acceptable range 88.9% (16 of 18)
<i>Percent of population <15 years of age living in states / prov. where both surveillance indicators are met</i> 53.8%	

Poliovirus History

The WPV1 viruses in Angola during the end of 2010 and in 2011 represent a continuing outbreak resulting from importation from India in 2007. Only one genetic cluster has been observed. In 2010, cases were distributed widely including the northeastern part of the country and included moderate regional transmission. The four cases from 2011 were located in the southern part of the country. The cases in 2011 represent continued circulation from 2010 and not a new importation, marking the fourth consecutive year of transmission of this virus. Based upon multiple isolates from 2010 with less genetic linkage than expected, there is indication that surveillance may have gaps at the sub-national level.

Data Completeness

Percent of NPAFP cases with unknown age	0%
Percent of 6-35 month old NPAFP cases with unknown dose history	8.9%